


This article was downloaded by:

On: 22 March 2010

Access details: *Access Details: Free Access*

Publisher *Informa Healthcare*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Critical Reviews in Biochemistry and Molecular Biology

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713609207>

### Cellular Dynamics of Small RNAs

Anita K. Hopper <sup>a</sup>

<sup>a</sup> Department of Biochemistry and Molecular Biology, Pennsylvania State University College of Medicine, Hershey, PA, USA

**To cite this Article** Hopper, Anita K.(2006) 'Cellular Dynamics of Small RNAs', *Critical Reviews in Biochemistry and Molecular Biology*, 41: 1, 3 – 19

**To link to this Article:** DOI: 10.1080/10409230500405237

**URL:** <http://dx.doi.org/10.1080/10409230500405237>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

# Cellular Dynamics of Small RNAs

## Anita K. Hopper

Department of Biochemistry  
and Molecular Biology,  
Pennsylvania State University  
College of Medicine, Hershey,  
PA, USA

**ABSTRACT** This review highlights the unexpectedly complicated nuclear egress and nuclear import of small RNAs. Although nucleus/cytoplasm trafficking was thought to be restricted to snRNAs of many, but not all, eukaryotes, recent data indicate that such traffic may be more common than previously appreciated. First, in conflict with numerous previous reports, new information indicates that *Saccharomyces cerevisiae* snRNAs may cycle between the nucleus and the cytoplasm. Second, recent studies also provide evidence that other small RNAs that function exclusively in the nucleus—the budding yeast telomerase RNA and possibly small nucleolar RNAs—may exit to the cytoplasm, only to return to the nucleus. Third, nucleus/cytoplasm cycling of RNAs also occurs for RNAs that function solely in the cytoplasm, as it has been discovered that cytoplasmic tRNAs of budding yeast travel “retrograde” to the nucleus and, perhaps, back again to the cytoplasm to function in protein synthesis. Fourth, there is at least one example in ciliates of small double-stranded RNAs traveling multiple cycles between the cytoplasm and distinct nuclei to direct genome structure. This report discusses data that support or argue against nucleus/cytoplasm bidirectional movement for each category of small RNA and the possible roles that such movement may serve.

**KEYWORDS** snRNA, snoRNA, telomerase RNA, TLC1, tRNA, dsRNA, siRNA, miRNA

## INTRODUCTION

Movement of macromolecules in and out of the nucleus occurs through aqueous channels that traverse the outer and inner nuclear membranes (ONM and INM, respectively). The channels are formed by nuclear pore complexes (NPCs), which join the ONM and INM. Both in lower and higher eukaryotes, NPCs are comprised of about 30 different proteins, called nucleoporins (Rout *et al.*, 2000; Cronshaw *et al.*, 2002). With the prominent exception of mRNAs (reviewed in Rodriguez *et al.*, 2004), transport of macromolecules through the NPC is generally mediated by a highly conserved mechanism consisting of the GTPase, Ran, Ran-regulatory proteins, and a family of Ran-binding proteins, the  $\beta$  importins. Members of the  $\beta$  importin family have the ability to specifically interact with particular cargoes and particular NPC proteins, moving with the cargo into or out of the nucleus. The  $\beta$  importin family is comprised of 14 members in budding yeast and ~20 in higher eukaryotes (reviewed in Gorlich & Kutay, 1999). The members are generally

Address correspondence to Anita K. Hopper, Department of Biochemistry and Molecular Biology, Pennsylvania State University College of Medicine, 500 University Drive, Hershey, PA 17033. E-mail: ahopper@psu.edu

referred to as importins if they move cargo from the cytoplasm to the nucleoplasm and as exportins if they move cargo from the nucleoplasm to the cytoplasm. However, it has been reported that some members can transport one specific cargo in one direction and another cargo in the opposite direction (*e.g.*, Yoshida & Blobel, 2001). Ran acts as a molecular switch regulating association or dissociation between a  $\beta$  importin family member and its cargo, dependent upon Ran's GTP or GDP bound state. The regulatory proteins determine whether Ran is associated with GTP (RanGEF) or GDP (RanGAP). Directionality across the nuclear pore is maintained by a gradient of RanGTP, which is high in the nucleus and low in the cytoplasm, due to the subcellular locations of RanGEF and RanGAP (Izaurrealde *et al.*, 1997; Kalab *et al.*, 2002). RanGEF is located in the nucleoplasm, whereas RanGAP is located primarily in the cytoplasm (Aebi *et al.*, 1990; Hopper *et al.*, 1990; Ohtsubo *et al.*, 1991).

There are numerous types of small RNAs encoded by the nuclear genome, and they serve a multitude of functions in various subcellular compartments. Some function strictly within the nucleus, whereas others function elsewhere in the cell. Nuclear genome-encoded small RNAs (snRNAs) U1, U2, U4, U5 and U6 and the corresponding AT/AC snRNAs function solely in the nucleus at various steps of pre-mRNA intron removal (reviewed in Tarn & Steitz, 1997). Likewise, a host of other small RNAs (snoRNAs) located in the nucleolar compartment of the nucleus function in pre-rRNA processing and modification (reviewed in Kiss, 2002). Another example of a small nucleus-encoded RNA with a strictly nuclear role is the small RNA component of telomerase (TLC1 in budding yeast), which functions in telomere maintenance and biogenesis (reviewed in Cech, 2004). In contrast, tRNAs provide an example of small RNAs that appear not to have a biological role in the nucleus, as they participate in protein synthesis in the cytoplasm. However, there has been a controversial report of translation in the nucleus (Iborra *et al.*, 2001; Review: Dahlberg *et al.*, 2003). If verified, it would mean that some part of the cellular tRNA pool functions in the nucleus. Furthermore, in some organisms, tRNAs encoded by the nucleus are imported into mitochondria and function in protein synthesis within this organelle (reviewed in Hopper & Phizicky, 2003). Finally, small double-stranded RNAs (dsRNAs) are the most recently discovered category of RNA. Some members of this category function in the cytoplasm in post-

transcriptional regulation of mRNA turnover and/or translation (reviewed in Tang, 2005), whereas other members function in the nucleus and regulate transcription and/or chromatin structure (reviewed in Matzke & Birchler, 2005; Bernstein & Allis, 2005). Whether a given dsRNA can function both in the nucleus and the cytoplasm is unknown.

As snRNA, snoRNA, and telomerase RNA are transcribed and function in the nucleus, one might have expected that they would never leave the nuclear compartment. Likewise, one would expect that tRNA movement in cells would be unidirectional—from the nucleus to the cytoplasm—because tRNAs are transcribed in the nucleus but function in the cytoplasm. However, in contradiction to the predictions, it has been established that U1, U2, U4, and U5 snRNAs have complicated cellular dynamics; they are exported to the cytoplasm where they undergo modification and associate with proteins, only to be re-imported back to the nucleus (Reviews: Kiss, 2004; Yong *et al.*, 2004). Surprisingly, complex nucleus/cytoplasm dynamics may not be restricted to snRNAs, as recent data indicate that other RNAs that function strictly in the nucleus may be exported to the cytoplasm and re-imported back into the nucleus. Moreover, it has been discovered recently that tRNAs in the cytoplasm can be imported into the nucleus—a process called “retrograde tRNA nuclear import” (Shaheen & Hopper, 2005; Takano *et al.*, 2005).

The focus here is threefold: (1) to assess the data indicating that small RNAs other than snRNAs that function exclusively in the nucleus may have a cytoplasmic phase; (2) to review the new developments documenting that resident cytoplasmic RNAs can be imported retrograde into the nucleus; and (3) to describe the nuclear/cytoplasmic dynamics of small dsRNAs; at least one member of this category of RNAs exhibits multiple cycles of nucleus egress and import. The review is organized to first discuss RNAs that function solely in the nucleus, followed by a discussion of RNA whose function is in the cytoplasm, and then by the dsRNA category of RNAs, some members of which function in the cytoplasm and some in the nucleus. Since the data in many of the recent publications rely upon a new application of yeast heterokaryon analysis, this methodology and its power/limitations are also assessed. Not discussed here are the RNAs that comprise the signal recognition particle, RNaseP, and U7 RNP, as there are limited published data concerning their movement among subcellular compartments. Also not elaborated upon

here are the complex intranuclear dynamics of resident nuclear RNAs; this subject has been thoroughly covered in recent reviews (Kiss, 2004; Lamond & Spector, 2003; Matera, 1999; Maser & DePinho, 2002).


## snRNAs—PRECEDENT FOR A CYTOPLASMIC PHASE FOR RESIDENT NUCLEAR RNAs

### Trafficking of U1, U2, U4 and U5 Between the Nucleus and the Cytoplasm

Cycling of U1, U2, U4, and U5 snRNAs between the nucleus and the cytoplasm is well established and extensively reviewed (reviewed in Kiss, 2004; Yong *et al.*, 2004). To set the stage for discussion of whether other RNAs also cycle between the nucleus and the cytoplasm, only the salient features of the complicated intracellular travels of U1, U2, U4, and U5 snRNAs are summarized (Figure 1).

U1, U2, U4, and U5 snRNAs are transcribed by RNA polymerase II (Pol II) into pre-snRNAs with extended 3' termini. First, these pre-snRNAs

acquire a 5' monomethyl guanosine cap ( $m^7GpppG$ , " $m^7G$ ") and then the nuclear cap-binding complex (CBC) interacts with  $m^7G$ -modified snRNAs (Hamm & Mattaj, 1990; Izaurralde *et al.*, 1995). Subsequently, the CBC/snRNA ribonucleoprotein (RNP) complex interacts with PHAX, which acts as an adaptor for the cooperative addition of the exportin, CRM1, and RanGTP, both required for nuclear export (Ohno *et al.*, 2000). Once in the cytoplasm, hydrolysis of RanGTP to RanGDP and dephosphorylation of PHAX result in the release of the snRNAs (Ohno *et al.*, 2000). The SMN protein complex facilitates assembly of the RNA-binding Sm proteins to the cytoplasmic snRNAs (Reviewed in Terns & Terns, 2001; Yong *et al.*, 2004), followed by hypermodification of  $m^7G$  to  $m^{2,2,7}G$  ("TMG") catalyzed by Tgs1 (Hausmann & Shuman, 2005) and 3' end maturation. The TMG moiety interacts with snurportin-1 (SPN1), which acts as an adaptor for importin- $\beta$ , providing one of the two nuclear import pathways (Fischer & Lüthmann, 1990; Huber *et al.*, 1998). The second import pathway is independent of TMG, but depends upon the Sm and SMN complexes; it also employs importin- $\beta$  (Narayanan *et al.*, 2004).



**FIGURE 1** Intracellular movements of U1, U2, U4, and U5 snRNAs between the nucleus and the cytoplasm. These RNAs are transcribed and partially processed in the nucleus. The  $m^7G$  modified 5' end is recognized by the cap binding complex (CBC) and then the complex interacts with PHAX that acts as an adaptor for the exportin, CRM1. Once released in the cytoplasm the RNAs interact with the SMN complex followed by binding of the Sm proteins, 5' cap modification to TMG and 3' end processing. The modified snRNP associates with snoportin (SPN) and the resulting RNP is imported back into the nucleus by two importin- $\beta$  dependent processes. Gray shaded bar, unprocessed 3' terminus; small circles,  $m^7G$  (1 circle) or TMG (3 circles) 5' modification; crescent shaped symbols, the cap-binding complex and PHAX as indicated.

It has been proposed that the cytoplasmic phase for the snRNAs provides a proofreading step that prohibits nuclear accumulation of nonfunctional snRNAs and prevents Sm protein assembly on inappropriate RNAs (Yong *et al.*, 2004). If the cytoplasmic cycle for snRNAs serves an important proofreading function, then it should be conserved throughout eukaryotes. However, several lines of evidence have argued against a cytoplasmic phase for snRNPs in budding yeast. First, PHAX homologs are not apparent in the yeast genome; so, there is no candidate protein that would be involved in snRNA nuclear export (Cited in: Murphy *et al.*, 2004). Second, yeast also do not encode SPN1 and SMN homologs involved in snRNP nuclear import (Cited in: Murphy *et al.*, 2004; Yong *et al.*, 2004). Third, it has been reported that 5' TMG formation and 3' end trimming occur strictly in the nucleus in yeast, rather than in the cytoplasm as in higher eukaryotes (van Hoof *et al.*, 2000; Mouaikel *et al.*, 2002). Finally, snRNAs have not been detected in the cytoplasm in budding yeast (cited in Murphy *et al.*, 2004).

Despite the evidence against nucleus/cytoplasm bidirectional movement for snRNAs in budding yeast, recent reports indicate that yeast snRNAs may indeed cycle between the nucleus and the cytoplasm. First, the U1-binding protein, Prp40, which participates in pre-mRNA splicing, was shown to move between the nucleus and the cytoplasm. Furthermore, Prp40 possesses leucine-rich nuclear export motifs recognized by Crm1 that are necessary for its function in splicing (Murphy *et al.*, 2004). One interpretation of the data is that Prp40 functions to export U1 snRNA from the nucleus to the cytoplasm (Murphy *et al.*, 2004).

Second, Olson and Siliciano (2003) provided direct evidence that snRNAs in budding yeast do cycle between the nucleus and the cytoplasm. They used a *kar1*-based heterokaryon assay established for assessing protein nucleus/cytoplasm bidirectional movement in budding yeast (Flach *et al.*, 1994). The *kar1* mutant causes a defect in nuclear fusion following mating, resulting in mated heterokaryon zygotes that contain separate parental nuclei within a common cytoplasm. Because yeast have a closed mitosis, gene products encoded by one nucleus and located in that nucleus can gain entrance to second nucleus within the joint cytoplasm only if the gene products have the ability to exit the first nucleus. A large number of experiments have documented that DNA exchanges occur only rarely between the nuclei of heterokaryons and that soluble

karyophilic proteins in one nucleus do not migrate to a second nucleus in the absence of nuclear export information (Dutcher, 1981; Sigurdson *et al.*, 1981; Flach *et al.*, 1994; Shaheen & Hopper, unpublished data).

Olson and Siliciano (2003) generated functional snRNA variants specifically detectable by fluorescence *in situ* hybridization (FISH). These variant snRNAs were provided to a haploid nucleus via a low copy plasmid, and the resulting cells were mated to a *kar1* haploid strain to generate heterokaryons. The subcellular dynamics of the tagged snRNAs were assessed using a modified version of the heterokaryon assay. About 50% of the heterokaryons demonstrated equal amounts of the snRNAs in each nucleus within ~2 h post mating, indicating movement from the first to the second nucleus. The authors provided convincing evidence that exit from one nucleus and entry to the second is not an artifact of “leaking” nuclei, because the process depends upon a functional exportin, Crm1 (Olson & Siliciano, 2003). However, because all of the various snRNA and snoRNAs tested (see below for further discussion of snoRNAs) demonstrated similar nucleus/cytoplasmic bidirectional movement, the authors suggested that “caution must be taken when testing nuclear envelope shuttling in yeast heterokaryons.” Nevertheless, the ability of Prp40 and snRNAs to move between the nucleus and the cytoplasm in yeast haploids and heterokaryons leaves open the possibility that components of the yeast splicing machinery cycle between the nucleus and the cytoplasm, similarly to the higher eukaryotic counterparts. Further studies are necessary to support or refute this conclusion.

## U6 snRNA Does Not Cycle Between the Nucleus and the Cytoplasm?

In contrast to U1, U2, U4, and U5 snRNAs, U6 snRNA appears not to exit the nucleus. Unlike the other snRNAs, U6 is transcribed by RNA polymerase III (Pol III). Also, unlike the other snRNAs, U6 does not possess Sm protein binding sites or a TMG cap; rather it possesses a methyphosphoryl G cap (Singh & Reddy, 1989). Although U6 snRNA that is injected into the cytoplasm of *Xenopus* oocytes is imported into the nucleus by a Ran-dependent process (Vankan *et al.*, 1990; Michaud & Goldfarb, 1992), data document that U6 snRNA transcribed in the nucleus is not exported to the cytoplasm. In particular, a catalytically active variant missing nuclear import information fails to accumu-

late in the cytoplasm after being injected into *Xenopus* oocyte nuclei (Vankan *et al.*, 1990). Assuming that the mutation did not alter putative nuclear export information, the data indicate that U6 RNA has no nuclear export potential. In support of this finding, variant U6 RNAs gain nuclear export ability when supplied with *cis*-acting nuclear export motifs (Terns *et al.*, 1993). On the other hand, budding yeast U6 snRNA moved simultaneously with U1 and U2 snRNAs in heterokaryons, although there are no published data to indicate that U6 nuclear egress is a Ran-dependent process (Olson & Siliciano, 2003). It has been suggested that U6's nuclear import capability in higher eukaryotes may serve to restore U6 snRNA nuclear pools after mitotic nuclear breakdown (Vankan *et al.*, 1990); however, since U6 snRNA is also imported into the nucleus in yeast and yeast has a closed mitosis, U6 snRNA nuclear import likely serves a different function, at least in budding yeast.

## NO CYCLING OF NUCLEAR RESIDENT snoRNAs BETWEEN THE NUCLEUS AND THE CYTOPLASM?

Cells encode a plethora of small RNAs that are located in the nucleolus (snoRNAs) and function in pre-rRNA nucleolytic processing and nucleoside modifications. Curiously, this category of RNAs has different biogenesis pathways in different organisms. In plants, snoRNAs are transcribed by Pol III and U3 snoRNA possesses a  $\gamma$  monomethyl phosphate cap (Shimba *et al.*, 1992). In contrast, in vertebrates snoRNAs are transcribed by Pol II, either as individual transcripts or within introns of pre-mRNAs (Shimba *et al.*, 1992; Tycowski *et al.*, 1996), and pre-snoRNAs possess a m<sup>7</sup>G cap that is modified to TMG on mature snoRNAs.

Given the similarities of vertebrate snoRNA biogenesis to snRNA biogenesis, it was anticipated, and reported (see citations in Peculis, 2001), that snoRNAs cycle between the nucleus and the cytoplasm. However, because snoRNAs lack motifs that bind Sm proteins, and binding Sm proteins, is important for their nuclear import, snoRNA nucleus/cytoplasm cycling was reinvestigated. It was learned that, in contrast to snRNAs, U3, U8, and U14 snoRNAs expressed in *Xenopus* oocytes could not be detected in the cytoplasm, unless vast excess of snoRNAs were injected into the nucleus or the cells possessed defects in the Ran cycle (Terns & Dahlberg, 1994; Terns *et al.*, 1995; Cheng *et al.*, 1995). Furthermore, and also in contrast to snR-

NAs, snoRNAs acquired the TMG modification in the nucleus, rather than the cytoplasm (Terns & Dahlberg, 1994). Subsequent studies employing HeLa and Ref52 cell cultures confirmed the results (Boulon *et al.*, 2004; Watkins *et al.*, 2004). So, even though snoRNAs and U1, U2, U4, and U5 snRNAs have a similar biogenesis pathway, their subcellular trafficking was reported to differ.

Studies of the role of CRM1 in snoRNA biogenesis showed that its inhibition by Leptomycin B (LMB) had no effect upon the maturation of U3 snoRNA, whereas the same regime had the anticipated inhibitory effect upon U1, U2, U4, and U5 pre-snoRNA processing (Boulon *et al.*, 2004). Rather, CRM1 was important for subnuclear localization because snoRNAs failed to accumulate in the nucleolus when CRM1 was inhibited by LMB. How cells distinguish between m<sup>7</sup>G-modified snRNAs that are exported to the cytoplasm and m<sup>7</sup>G-modified snoRNAs that are apparently retained in the nucleus has been investigated. Boulon and coworkers (2004) and Watkins and associates (2004) showed that m<sup>7</sup>G modified pre-U3 snoRNAs interact with PHAX and this interaction directs the pre-snoRNAs to Cajal bodies. TMG modification and 3'-end modification occur in the Cajal bodies and precede interaction of the matured snoRNA with CRM1. Rather than exporting the RNA to the cytoplasm, CRM1 appears to direct snoRNAs to the nucleolus (Boulon *et al.*, 2004). Thus, surprisingly, the same machinery necessary for U1, U2, U4, and U5 snRNA nuclear export may function in intranuclear movement of m<sup>7</sup>G-modified snoRNAs.

Do the data for snoRNA biogenesis exclude a cytoplasmic phase for this category of RNAs? Previously, Cheng and colleagues (1995) showed that in the absence of a functional Ran cycle, part of the U3 snoRNA is cytoplasmic. Although an interpretation of the results is that RanGTP functions in U3 snoRNA nucleolar retention (Cheng *et al.*, 1995), a recent study provided no support for this finding (Narayanan *et al.*, 2003). Likewise, although the studies of the roles of PHAX and CRM1 in snoRNA biogenesis indicate that these proteins function in snoRNA intranuclear trafficking, both Boulon and colleagues (2004) and Watkins and coworkers (2004) considered the alternative possible interpretation that PHAX and CRM1 function in snoRNP nuclear export. If indeed all or part of the pool of this category of snoRNA exits the nucleus, then inhibition of the nuclear import machinery should cause their cytoplasmic accumulation. Employing RNAi methodology, Watkins and colleagues (2004)

reported that down-regulation of SMN causes reduced U3, but not U8 or U14, levels; however, the location of these snoRNAs upon RNAi treatment has not yet been published. Likewise, if snoRNAs cycle between the nucleus and the cytoplasm, snoRNAs introduced into the cytoplasm should be able to access the nuclear interior. It is known that U3 complex-associated proteins moves bidirectional between the nucleus and the cytoplasm (Leary *et al.*, 2004). However, studies of the capability of U3 snoRNA to be imported into the nucleus have led to contradictory conclusions. One report (Speckmann *et al.*, 1999) detected no significant import of U3 or U8 snoRNAs into *Xenopus* nuclei when these RNAs were injected into the cytoplasm, whereas others, also using the *Xenopus* system, reported that there is snoRNA nuclear import, albeit a lower rate and reduced efficiency compared to snRNAs (Baserga *et al.*, 1992; Peculis, 2001). It is possible that a nuclear import salvage pathway is able to retrieve snoRNAs that somehow escape the nuclear retention process (Terns *et al.*, 1995). Nevertheless, the question of whether U3, U8, and U14 snoRNAs cycle between the nucleus and the cytoplasm may warrant further investigation.

## POSSIBLE CYCLING OF THE NUCLEAR RESIDENT RNA SUBUNIT OF TELOMERASE BETWEEN THE NUCLEUS AND THE CYTOPLASM

Telomerase is a RNP consisting of a single RNA (TLC1 in budding yeast) and multiple proteins. The RNA subunit serves as the template and DNA synthesis is catalyzed by the reverse transcriptase subunit of the RNP complex, TERT/Est2 (Lingner *et al.*, 1997; reviewed in Cech, 2004). In yeast, the other two proteins of the enzyme complex, Est1 and Est3, are required only *in vivo* for appropriate maintenance of telomere length (Lendvay *et al.*, 1996; review: Cech, 2004). Yeast and vertebrate telomerase RNA are transcribed by Pol II, but the ciliate RNA is transcribed by Pol III (reviewed in Collins, 1999). The vertebrate telomerase RNA contains H/ACA motifs typically found in snoRNAs that function in snoRNA biogenesis and subnuclear distribution (Mitchell *et al.*, 1999; Narayanan *et al.*, 1999; Lukowiak *et al.*, 2001). Like for snoRNAs, the vertebrate telomerase RNA appears to never exit the nucleus (Lukowiak *et al.*, 2001); rather, the H/ACA motif has been implicated in telomerase RNA nuclear retention

(Lukowiak *et al.*, 2001). In contrast, budding yeast TLC1 possesses a TMG cap and associates with Sm proteins, so, its biogenesis resembles the biogenesis of snRNAs (Seto *et al.*, 1999). Since it has been thought that yeast snRNAs do not traffic between the nucleus and the cytoplasm, it was anticipated that the yeast telomerase RNA also would not exit the nucleus. However, two reports support the notion that budding yeast TLC1 may move between the nucleus and the cytoplasm.

Teixeira and colleagues (2002) employed the yeast heterokaryon assay to study the intracellular trafficking of TLC1. Because FISH technology is not sensitive enough to detect endogenous levels of TLC1, the authors used a plasmid encoding a *GAL1*-induced, over-expressed TLC1. Haploid cells containing the inducible *TLC1* gene were mated to haploid cells harboring a *kar1* mutation. For 10% to 60% of the heterokaryons, TLC1 RNA was present in all heterokaryon nuclei and not just the nucleus that contained the plasmid encoding TLC1 RNA. The data indicate that TLC1 can exit one nucleus and gain entrance to a second nucleus through the shared cytoplasm.


Two caveats complicate interpretation of the data: (1) the studies tracked considerably over-expressed RNA, and (2) the location of TLC1 was assessed 12 h. after heterokaryon formation. *A priori*, it is possible that over-expressed TLC1 could overwhelm a putative nuclear retention mechanism and diffuse out of the nucleus. However, unlike the ciliate or vertebrate telomerase RNAs that are quite small (150 and 450 nt., respectively), TLC1 from budding yeast is large (1.2 kb, ~400 kd) and very unlikely to exit the nucleus via diffusion because it greatly exceeds the size of the nuclear pore diffusion channel (<47 kd; Shulga *et al.*, 2000). Nevertheless, to address the caveat of over-production Teixeira and coworkers (2002) also generated a construct in which the TLC1 transcript was expressed in endogenous levels but was fused to another RNA encoding an ORF. This fusion transcript was translated and generated the protein encoded by the ORF. Since translation occurs only in the cytoplasm, production of the ORF protein required that fusion transcript to be exported to cytoplasm. One interpretation of the data is that TLC1 nuclear export can occur in the absence of its over-expression. However, the authors were unable to eliminate the alternative that the ORF-encoding part of the fusion transcript was responsible for directing the RNA to the cytoplasm. The second caveat concerns the interval between heterokaryon formation and FISH analysis,

as this length of time could be sufficient for TLC1 RNA that managed to “escape” from the nuclear interior to interact with newly synthesized TLC1-binding proteins and be imported in tow as RNPs. However, no cytoplasmic TLC1 pools were evident, indicating a rather efficient nuclear import process.

Nucleus/cytoplasmic bidirectional movement for *S. cerevisiae* TLC1 is supported by an independent study. Employing a synthetic genetic interaction screen for yeast mutants defective in telomere maintenance, Ferrezuelo and colleagues (2002) identified *mtr10*. Biochemical studies showed that *mtr10* mutant cells have short telomeres. A deletion of *MTR10* also resulted in low cellular levels of telomerase, similar to cells with defects in the telomerase protein subunits, Est1, Est2, and Est3. TLC1 processing was also altered in *mtr10*Δ cells. *MTR10* encodes 1 of the 14 members of the β importin family of Ran-binding proteins and it has a well established function in nuclear import of mRNA binding proteins (e.g., Senger *et al.*, 1998). The authors eliminated the possibility that the phenotype of the *mtr10*Δ mutant cells was due to a defect in nuclear import of one of the known Mtr10 substrates, because mutation of the substrate did not cause defects in telomere metabolism. The authors also eliminated the possibility that low levels of telomerase resulted from defects in nuclear import of Sm proteins because *mtr10*Δ had normal levels of snRNAs that require nuclear import of Sm proteins for their stability. FISH analysis

was employed to determine whether *mtr10*Δ altered the subcellular distribution of TLC1. Interestingly, whereas U3 snoRNA was located in the nucleus in both wild-type and *mtr10*Δ cells, *GAL1*-regulated over-expressed TLC1 was mislocated to the cytoplasm in *mtr10*Δ, but not in wild-type cells. The simplest interpretation of why TLC1 is mislocated to the cytoplasm is that it normally moves bidirectionally between the nucleus and the cytoplasm and that Mtr10 functions in its nuclear import. However, the authors were unable to eliminate the alternative interpretation that Mtr10 is responsible for import of a product needed for processing of TLC1 in the nucleus and in the absence of this processing step, TLC1 aberrantly exits the nucleus. Also, since in these experiments TLC1 was also over-expressed, it is possible that “escaped” TLC1 RNA can be re-imported as an RNP with newly synthesized Est proteins and that import of the RNP is an Mtr10-mediated process.

In combination, the heterokaryon studies that document TLC1 movement from one nucleus to another (Teixeira *et al.*, 2002), the ability of TLC1-ORF fusion transcripts to be translated (Teixeira *et al.*, 2002), and the location of TLC1 in the cytoplasm in *mtr10*Δ cells (Ferrezuelo *et al.*, 2002), provide tantalizing evidence that TLC1 traffics between the nucleus and the cytoplasm, at least in budding yeast (Figure 2). If this RNA indeed moves bidirectionally in and out of the nucleus, it will be important to discover its significance and to characterize mechanism responsible for its export to the



**FIGURE 2** Possible nucleus/cytoplasm dynamics for the yeast telomerase RNA subunit—TLC1. The mechanism of putative export to the cytoplasm is unknown. The importin Mtr10 is proposed to transport the RNA back into the nucleus. 3' and possibly 5', modifications may take place after the RNA returns to the nucleus. Symbols are as in Figure 1.

cytoplasm. Along these lines, Boulon and colleagues (2004) demonstrated physical interaction of PHAX and telomerase RNA in HeLa cell extracts. As detailed above PHAX interaction with snRNAs mediates their Ran-dependent nuclear export; however, because PHAX interaction with U3, U8, and U13 snoRNA may function in their intranuclear dynamics rather than their nuclear export, the interactions are difficult to interpret. Future studies in yeast and higher eukaryotes are necessary to resolve whether telomerase RNA does indeed cycle between the nucleus and the cytoplasm.

## **tRNA NUCLEAR EXPORT AND IMPORT—PRECEDENT FOR NUCLEAR IMPORT FOR A RESIDENT CYTOPLASMIC RNA**

Cytoplasmic tRNAs are transcribed in the nucleus where they undergo numerous post-transcriptional alterations. The tRNAs then exit the nucleus and undergo further modification and processing in the cytoplasm (reviewed in Hopper & Phizicky, 2003). Completely processed tRNAs acquire appropriate amino acids at their 3' ends, catalyzed by 1 of 20 aminoacyl-tRNA synthetases, and the charged tRNAs deliver the amino acids to ribosomes where amino acids are assembled into proteins at the direction of mRNA codons. Given this simple "central dogma" view of the role of tRNA in gene expression, one would anticipate that the movement of tRNAs in cells is unidirectional, nucleus to cytoplasm, and that tRNA aminoacylation would be restricted to the cytoplasm. Both of these predictions are incorrect. Moreover, the subcellular organization of the tRNA biogenesis pathway is unexpectedly complex.

## **Complex Intranuclear and Cytoplasmic Organization of the tRNA Biogenesis Machinery**

As transcribed, pre-tRNAs are oversized on the 5' and 3' termini and may contain an intron. Conversely, tRNA transcripts lack the numerous nucleoside modifications (>47 known nucleoside modifications in eukaryotes, 25 in *S. cerevisiae*; Rozenski *et al.*, 1999; Hopper & Phizicky, 2003) that reside on mature tRNAs, as well as the 3' CCA sequence that is required for aminoacylation (reviewed in Hopper & Phizicky, 2003). Moreover, one tRNA, tRNA<sup>His</sup>, acquires a post-transcriptional 5' additional G (Gu *et al.*, 2003). In budding yeast at least

60 gene products, about 1% of the genome, are required to convert pre-tRNAs to mature functional molecules. In all eukaryotes many of the processing steps occur before tRNAs exit the nucleus to the cytoplasm. However, steps that occur in the nucleus in one organism may occur in the cytoplasm of another. For example, pre-tRNA splicing is clearly nuclear in vertebrates (De Robertis *et al.*, 1981; Lund & Dahlberg, 1998; Paushkin *et al.*, 2004), while this process occurs in the cytoplasm, on the surface of mitochondria, in yeast (Huh *et al.*, 2003; Yoshihisa *et al.*, 2003; Shaheen & Hopper, 2005).


The intranuclear locations of the tRNA transcription and processing machinery is highly organized in *S. cerevisiae*. Despite the fact that there are ~275 tRNA-encoding genes dispersed throughout the genome, their transcription is localized to the nucleolus (Thompson *et al.*, 2003; reviewed in Haeusler & Engelke, 2004). How this is achieved topologically is unknown, but it does put significant three-dimensional constraints on the genome. Removal of the 5' leader sequence, catalyzed by RNase P, a 10-subunit RNP complex, also occurs in the nucleolus in yeast (Bertrand *et al.*, 1998), but perhaps not in vertebrate cells (reviewed in Xiao *et al.*, 2002). In contrast, for 19 proteins known to catalyze tRNA modifications steps in the nucleus (Huh *et al.*, 2003; Hopper & Phizicky, 2003), only one, Mod5, appears to be concentrated in the nucleolus (Tolerico *et al.*, 1999). Trm1 (Rose *et al.*, 1992; Rose *et al.*, 1995) and possibly Trm4 (Wu *et al.*, 1998) reside at the inner nuclear membrane, and the remaining 16 to 17 tRNA modification proteins appear to be dispersed throughout the nucleoplasm (Huh *et al.*, 2003). Nine, or possibly 10, of these proteins also have cytoplasmic pools, whereas 6, or possibly 7, modification proteins reside strictly in the cytoplasm. Cytoplasmic tRNA processing activities are also specifically located. For example, the four splicing endonuclease complex subunits are located on the cytoplasmic surface of mitochondria, whereas the enzymes that ligate the two tRNA halves together and remove the residual 2' phosphate from the splice junction are located throughout the cytoplasm (Yoshihisa *et al.*, 2003; Huh *et al.*, 2003). Thus, for *S. cerevisiae*, tRNA processing activities reside in numerous subnuclear compartments—the nucleolus, the nucleoplasm, and the inner nuclear membrane—as well as in particular domains of the cytoplasm. What function(s) this elaborate organization serves and whether similar organizational patterns also exist in higher eukaryotes, remains to be discovered.

## Parallel Pathways for tRNA Egress to the Cytoplasm

There is one known exportin (Los1 in yeast, Exportin-t [Xpo-t] in vertebrates, and PSD in plants) dedicated to transporting tRNA to the cytoplasm (Arts *et al.*, 1998a; Hellmuth *et al.*, 1998; Kutay *et al.*, 1998; Sarkar & Hopper, 1998; Hunter *et al.*, 2003; Park *et al.*, 2005). Xpo-t binds tRNA in the nucleus in the presence of RanGTP and it is able to interact with two distinct nucleopore proteins (Kuersten *et al.*, 2002). Vertebrate Xpo-t likely also serves a proofreading function because it preferentially interacts with tRNAs with correct tertiary structure and completely processed 5' and 3' termini (Arts *et al.*, 1998b; Lipowsky *et al.*, 1999). Interestingly, this exportin does not distinguish between intron-containing and spliced tRNAs (Arts *et al.*, 1998b; Lipowsky *et al.*, 1999). Upon movement of the heterotrimeric tRNA, RanGTP, and Xpo-t complex to the cytoplasm, RanGAP stimulates GTP hydrolysis, and the tRNA cargo is released from the complex into the cytoplasm (Figure 3).

Despite the simplicity and elegance of this nucleus-to-cytoplasm delivery mechanism, it fails to account for the egress of most tRNA to the cytosol for several organisms. As first demonstrated in yeast, the gene en-

coding this exportin, *LOS1*, is unessential (Hurt *et al.*, 1987); in fact, *los1Δ* cells are quite hardy. *LOS1* homologs also have been shown to be unessential in those organisms, *S. pombe* and *Arabidopsis*, in which the gene has been ablated (Sazer, personal communication; Li & Chen, 2003), implicating an alternative Los1-independent tRNA nuclear egress pathway(s). Although no genetic studies of the necessity of the vertebrate Xpo-t have been reported, it has been thought to be responsible for the export of the vast majority of tRNA from the nucleus to the cytoplasm because its inhibition via antibody injection causes nuclear retention of nearly the entire tRNA pool (Arts *et al.*, 1998b; Lipowsky *et al.*, 1999). Nevertheless, vertebrates encode another exportin, Exp5, that is able to bind tRNA (Bohnsack *et al.*, 2002; Calado *et al.*, 2002) and may serve to export tRNA from the nucleus (Calado *et al.*, 2002). It has been reported that yeast possessing deletions of both the Xpo-t and the Exp5 homologs, Los1 and Msn5, respectively, accumulate more tRNA in the nucleus than either alone, perhaps implicating Msn5 in tRNA nucleus/cytoplasm dynamics in yeast (Takano *et al.*, 2005). However, because *msn5Δ* cells are reported not to accumulate tRNA in the nucleus, and *los1Δ msn5Δ* double mutants have no synergistic growth defects (Feng & Hopper, 2002; Takano *et al.*, 2005) it would appear



**FIGURE 3** tRNA in *S. cerevisiae* moves retrograde from the cytoplasm to the nucleus. Pre-tRNAs interact directly with the exportin, Los1, in the nucleus and are released into the cytoplasm. Further modification, including removal of introns occurs in the cytoplasm. Retrograde movement of the mature tRNAs to the nucleus under nutrient deprivation depends upon the importin, Mtr10. Likely, tRNAs are exported a second time to the cytoplasm under favorable growth conditions. Gray shaded bar = intron; small circle = amino acid.

that Msn5 does not provide a major export pathway in budding yeast. Moreover, because Exp5 interacts more poorly with tRNA than does Xpo-t, it is also thought to provide only a minor nuclear export pathway for tRNA in vertebrates (Calado *et al.*, 2002). Consistent with this idea, it has been shown that deletion of the Exp5 homolog, HST, from the *Arabidopsis* genome causes no defect in tRNA nuclear export (Park *et al.*, 2005). Rather, Exp5 and HST bind dsRNAs, such as precursors to micro-RNAs (pre-miRNA), and serve as their nuclear export receptor (Lund *et al.*, 2004; Zeng & Cullen, 2005; Gwizdek *et al.*, 2004; Park *et al.*, 2005). Since the combination of *los1* $\Delta$ /PSD and *msn5* $\Delta$ /HST is not lethal in yeast or *Arabidopsis* (Takano *et al.*, 2005; Park *et al.*, 2005), the genetic data indicate the existence of additional unaccounted tRNA nuclear export pathways, at least in these organisms.

## A Role for tRNA Aminoacylation in tRNA Nuclear Egress

Contrary to dogma, Lund & Dahlberg (1998) showed that tRNAs can be aminoacylated in the nuclear interior of *Xenopus* oocytes. Both genetic and biochemical studies showed that nuclear tRNA aminoacylation also occurs in budding yeast (Sarkar *et al.*, 1999; Steiner-Mosonyi & Mangroo, 2004). In vertebrates, nuclear tRNA aminoacylation is important, although not essential, for tRNA nuclear export (Lund & Dahlberg, 1998; Arts *et al.*, 1998b). Conditional mutations of genes encoding tyrosyl-tRNA synthetase, isoleucyl-tRNA synthetase, and methionyl-tRNA synthetase, each cause tRNA nuclear accumulation (Sarkar *et al.*, 1999), and inhibitors of isoleucyl-tRNA synthetase likewise cause tRNA nuclear accumulation (Grosshans *et al.*, 2000), indicating that nuclear tRNA aminoacylation is also important for tRNA nuclear export in budding yeast. Consistent with this finding, blocking the addition of the C, C, and A nucleotides that are requisite for aminoacylation or depletion of amino acids from growth media can cause tRNA nuclear accumulation (Sarkar *et al.*, 1999; Grosshans *et al.*, 2000; Feng & Hopper, 2002; Shaheen & Hopper, 2005; Takano *et al.*, 2005).

If indeed tRNAs can be aminoacylated in the nucleus as well as in the cytosol, then there should be nuclear pools of the aminoacyl-tRNA synthetases. It has been shown by both microscopic analyses and subcellular fraction studies that there are small pools of aminoacyl-tRNA synthetases in vertebrate nuclei (Popenko *et al.*,

1994; Nathanson & Deutscher, 2000). Likewise, a small pool of tyrosyl-tRNA synthetase, Tys1, co-fractionates with budding yeast nuclei (Azad *et al.*, 2001). Moreover, Tys1 possesses a functional nuclear localization sequence (NLS) that quantitatively delivered a passenger  $\beta$ -galactosidase to the nucleus (Azad *et al.*, 2001). Mutations of the NLS reduced the pool of Tys1 that co-fractionated with nuclei, but caused no significant defect in Tys1 catalytic activity. The NLS mutations resulted in increased nuclear tRNA pools, consistent with the hypothesis that nuclear pools of aminoacyl-tRNA synthetases are important for tRNA nucleus/cytoplasm distribution (Azad *et al.*, 2001).

## Surprising Retrograde Traffic of tRNA from the Cytoplasm to the Nucleus

In vertebrate cells, pre-tRNA splicing occurs in the nucleus (De Robertis *et al.*, 1981; Lund & Dahlberg, 1998; Paushkin *et al.*, 2004), prior to tRNA end maturation (Lund & Dahlberg, 1998). In contrast, in budding yeast and, perhaps in *Arabidopsis*, splicing follows end processing (O'Connor & Peebles, 1991; Hiley *et al.*, 2005; Park *et al.*, 2005). Alteration of components of the Ran-dependent tRNA nuclear export pathway (RanGAP, RanGEF, or Los1p), as well as deletions of numerous nucleoporins, each cause accumulation of end-processed, intron-containing pre-tRNAs (Hopper *et al.*, 1978; Hopper *et al.*, 1980; Kadowaki *et al.*, 1994; Sharma *et al.*, 1996). The discovery that pre-tRNA splicing occurs in the cytoplasm in budding yeast (Yoshihisa *et al.*, 2003) provided an explanation for the defects in pre-tRNA intron removal that accompany alterations of the tRNA nuclear export machinery as pre-tRNAs maintained in the nucleus would be unable to access the cytosolic splicing endonuclease. However, not all mutations or physiological conditions that cause nuclear accumulation of tRNA in yeast result in defects in pre-tRNA intron removal. For example, inhibitors of aminoacylation, conditional mutations of aminoacyl-tRNA synthetases, Cca1, and the translation elongation factor EF1, or amino acid deprivation, each cause nuclear accumulation of spliced tRNA (Grosshans *et al.*, 2000; Feng & Hopper, 2002; Shaheen & Hopper, 2005). Splicing of pre-tRNAs in the cytoplasm does not provide an explanation for the presence of nuclear pools of spliced, mature tRNAs. If tRNA transport is unidirectional—nucleus to cytoplasm—nuclear accumulation of spliced tRNA is in

conflict with splicing occurring solely in the cytoplasm. Two possible scenarios could explain this enigma: either (1) the splicing machinery or (2) mature, spliced tRNAs move retrograde from the cytoplasm to the nucleus.

The subcellular location of a tagged version of a catalytic subunit of splicing endonuclease, Sen2-GFP (Huh *et al.*, 2003) in cells grown in rich media was compared to its location in cells depleted for amino acids or cells with *tys1-1*—both causing nuclear pools of spliced tRNAs (Sarkar *et al.*, 1999; Grosshans *et al.*, 2000; Azad *et al.*, 2001; Whitney *et al.*, in preparation). Under all conditions, detectable Sen2-GFP was associated with mitochondria, arguing against the first scenario that the pre-tRNA splicing complex is able to move from the cytoplasm to the nucleus (Shaheen & Hopper, 2005).

Two groups employed heterokaryon analysis to investigate the unorthodox alternative that nuclear pools result from retrograde movement of spliced tRNAs from the cytosol to the nucleus. Shaheen and Hopper (2005) provided one of the mating haploids with a gene expressed from a plasmid maintained in a single copy in yeast and regulated by its endogenous promoter from *D. discoideum*. This gene encodes an intronless suppressor tRNA, tRNA<sup>Glu-D</sup>, known to participate in translation in yeast (Dingermann *et al.*, 1992). Takano and colleagues (2005) employed a tandem gene pair from *S. pombe* encoding intron-containing tRNA<sup>Ser</sup><sub>UGA</sub> and intronless tRNA<sup>Met</sup><sub>i</sub>. Because tRNA is normally efficiently exported from the nucleus, and because efficient export would ablate detection of tRNA imported into the nucleus, heterokaryon analyses were conducted using mutations (*los1Δ* or *los1Δ msn5Δ*) or physiological conditions (amino acid deprivation) known to cause tRNA nuclear accumulation. If tRNA moves retrograde to the nucleus, then tRNA encoded by only one nucleus of a heterokaryon would be detected in both nuclei. In contrast, if tRNA movement is unidirectional, *i.e.*, strictly nucleus to the cytosol, tRNA would not be detected in the second nucleus. As assessed by FISH, tRNA signals were detected in both nuclei of the heterokaryons, supporting the hypothesis that tRNA encoded by one nucleus can gain access to a second nucleus and providing evidence for retrograde movement of tRNA from the cytoplasm to the nucleus (Shaheen & Hopper, 2005; Takano *et al.*, 2005).

Both groups also demonstrated retrograde movement of endogenous tRNAs in haploid yeast. Shaheen and Hopper (2005) took advantage of the fact that

amino acid deprivation caused tRNA retrograde movement in the heterokaryon assay, and showed that endogenous tRNAs accumulate in the nucleus when wild-type haploid cells were deprived for amino acids. Takano and coworkers (2005) studied retrograde tRNA movement in cells in rich media, using a regimen established for assessing nuclear envelope permeability (Shulga *et al.*, 2000). This assay employs 2-deoxyglucose and sodium azide to equilibrate macromolecules across the nuclear membrane. Cells with defects in tRNA nuclear export (*los1Δ msn5Δ*) were treated with 2-deoxyglucose and sodium azide and then the location of cellular tRNA was monitored in re-fed cells in the absence of new tRNA synthesis. Transcription of new tRNAs was inhibited by thiolutin, a general inhibitor of all three RNA polymerases. tRNA accumulated in nuclei in the absence of new transcription, providing support for retrograde import from the cytoplasm (Takano *et al.*, 2005). Thus, the two groups provided evidence to support the paradigm-shifting idea that tRNAs move retrograde from the cytoplasm to the nucleus (Figure 3).

The results from both groups are in agreement that tRNA movement into nuclei is an active process. However, the role of the Ran cycle in tRNA retrograde movement to the nucleus is somewhat controversial. Nuclear accumulation of cytoplasmic tRNAs resulting from amino acid deprivation appeared to require a functional Ran cycle and the Mtr10 member of the  $\beta$  importin family (Shaheen & Hopper, 2005). However, a functional Ran cycle was not necessary for tRNA nuclear accumulation following 2-deoxyglucose, sodium azide, and thiolutin treatment (Takano *et al.*, 2005). Assuming both types of experiments faithfully report the retrograde process, it is possible that a functional Ran cycle is needed only for retrograde movement signaled by amino acid deprivation (Figure 3). The possible role of Mtr10 in tRNA nuclear import is interesting in light of its implicated role in nucleus/cytoplasm dynamics of telomerase RNA discussed above.

The ability of tRNA to enter the nucleus—“retrograde tRNA nuclear import”—reveals previously unknown complexity in its nucleus/cytoplasm dynamics. The purpose that this movement serves to cells is unknown. However, based upon the hypothesis that snRNAs move to the cytosol and then back to the nucleus so that they may be proofread in a different cellular compartment than that in which they function (Yong *et al.*, 2004), tRNA retrograde movement may likewise provide a proofreading role for tRNA processing steps

that occur in the cytoplasm, thereby preventing interaction of faulty tRNAs with the translation machinery (Shaheen & Hopper, 2005; Takano *et al.*, 2005). Alternatively, based upon the movement of mRNAs from ribosomes to P-bodies under nutrient deprivation (Teixeira *et al.*, 2005), retrograde tRNA nuclear import may serve a regulatory role by separating tRNAs from the translation machinery when active translation can not occur (Shaheen & Hopper, 2005). A model consistent with the evidence that aminoacylation affects tRNA nuclear export (Lund & Dahlberg, 1998; Sarkar *et al.*, 1999; Grosshans *et al.*, 2000; Azad *et al.*, 2001) suggests that tRNA aminoacylation regulates both tRNA retrograde import and re-export to the cytoplasm such that, upon appropriate signaling events, cytoplasmic tRNAs are delivered into the nucleus and are maintained there until the nutritional status is sufficient for nuclear tRNA aminoacylation can occur, providing a nutritional checkpoint prior to a second round of tRNA nuclear export (Figure 3).


Several burning questions remain to be answered: (1) Is tRNA retrograde movement constitutive and/or regulated? (2) What gene products are involved in the import and the putative nuclear retention processes? (3) Are tRNAs imported to the nucleus returned to the cytoplasm for translation or, instead, are they turned over by the nucleus-located tRNA degradation machinery (Kadaba *et al.*, 2004)? (4) Does tRNA retrograde movement occur in vertebrate cells? Since pre-tRNA splicing occurs in the nucleus in vertebrate cells, tRNAs may be proofread before leaving the nucleus, eliminating the need for import and a second round of export to assure that only mature functional tRNAs are in the same compartment with the translation machinery. On the other hand, since there is evidence for yeast that nuclear accumulation of cytoplasmic tRNAs responds to nutrient cues, tRNA retrograde movement might provide a new level of regulation of gene expression in response to nutritional deprivation for all eukaryotes.

## SMALL DOUBLE-STRANDED RNAs FUNCTION IN THE CYTOPLASM AND THE NUCLEUS

Since the discoveries of small double-stranded RNAs (dsRNAs) in the 1990s, these RNAs have been implicated in numerous processes, including mRNA turnover, mRNA translation, regulation of transcrip-

tion, chromatin structure and silencing, and genome stability. There are two categories of endogenous small dsRNAs: (1) small interfering RNAs (siRNA) and (2) micro RNAs (miRNA). Endogenous siRNAs are generally encoded by two complementary RNAs that can perfectly base pair with their targets, whereas miRNAs are encoded by self-complementary hairpin structures able to form interrupted base pairs with their targets. RNA-induced silencing complexes (RISC) use one strand of the dsRNAs to inactivate particular mRNAs by complementary base pairing, either by targeting the mRNAs for turnover or by inhibiting target mRNA translation, likely within cytoplasmic P-bodies (Bagga *et al.*, 2005; Liu *et al.*, 2005; Pillai *et al.*, 2005; Sen & Blau, 2005). Other siRNAs assemble into RNA-induced silencing complexes (RITS) that are recruited to chromatin and function in transcription silencing, and/or genome stability. The biogenesis (reviewed in Tomari & Zamore, 2005; Sontheimer, 2005) and biological roles of small dsRNAs (reviewed in Bartel, 2004; Pasquinelli *et al.*, 2005; Bernstein & Allis, 2005) have been extensively reviewed. Here, only their nucleus/cytoplasmic dynamics are explored.

miRNAs are encoded in the nucleus, but generally function in the cytoplasm. The mechanism for their nuclear egress has recently been described. In vertebrates the genes encoding miRNAs are transcribed into self-complementary hairpin initial precursor RNAs (pri-miRNAs) that undergo a two-step maturation process. In the nucleus, pri-miRNAs are processed to ~70 nucleotide double-stranded hairpin precursor miRNAs (pre-miRNAs) catalyzed by an RNase III-like enzyme, Drosha (Lee *et al.*, 2003). Pre-miRNAs directly interact with Exp5, an exportin member of the  $\beta$  importin family, in the presence of RanGTP. Subsequently, the heterotrimeric complex is exported from the nucleus to the cytoplasm (Lund *et al.*, 2004; Yi *et al.*, 2003; Bohnsack *et al.*, 2004). In the cytoplasm pre-miRNAs are processed to ~22-nucleotide double-stranded mature and functional microRNAs (miRNAs) by Dicer, another member of the RNase III category of endonucleases (Figure 4, left). This two-step biogenesis process involving two separate subcellular compartments appears to be conserved because there are Drosha homologs in mammals, *Xenopus*, *C. elegans*, and *Drosophila*, although not in *Arabidopsis* or *S. pombe* (Review: Murchison & Hannon, 2004). Nevertheless, in *Arabidopsis*, the steady state level of most, but not all, miRNAs is dependent upon the Exp5 homolog, *HASTY* (*HST*) (Park *et al.*,



**FIGURE 4** Intracellular dynamics of dsRNAs. Left: miRNAs. Initial precursor miRNAs are processed by Drosha to ~70 nucleotide hairpins. The resulting pre-miRNAs interact directly with the exportin, Exp5, and are transported to the cytoplasm. In the cytoplasm, they are processed further to 21-nucleotide mature dsRNA by Dicer before one strand enters the RISC complex. Right: scnRNA, an example of siRNAs that undergo several cycles between the nucleus and the cytoplasm. ScnRNAs generated by action of a Dicer-like enzyme in the nucleus are transported to the cytoplasm by an unknown mechanism. The RNAs are imported to the old macronucleus, also by an unknown mechanism, where they scan the genome for deleted DNA sequences; RNAs with no homolog are exported to the cytoplasm and imported to the new macronucleus where they specify deletion of the appropriate genomic sequences.

2005), perhaps indicative of nuclear export, despite the absence of Drosha.

Some siRNAs are recruited to sites on chromatin (Noma *et al.*, 2004; Reviewed in Martienssen *et al.*, 2005; Bernstein & Allis, 2005) where they may guide DNA rearrangements and/or transcriptional silencing, processes requiring transcription of the target DNA by Pol II (Schramke *et al.*, 2005; Kato *et al.*, 2005). Whether the siRNAs that function in the nucleus are first processed in the nucleus and then in the cytoplasm, like miRNAs that assemble into RISC complexes, or whether they are solely processed in the nucleus and assemble into RITS complexes in the nucleus, is not completely resolved. However, since deletion of the gene encoding the HST exportin from *Arabidopsis* appears to have no effect upon the steady state level of several tested endogenous siRNAs involved in post-transcriptional gene silencing (Park *et al.*, 2005), siRNA nuclear export and re-import may not occur in plants. In contrast, there is at least one example in *Tetrahymena* where special-

ized siRNAs (siRNA-like scan, scnRNAs) are thought to travel from the original micronucleus to the cytoplasm, to a second nucleus—the old macronucleus, and eventually to a third nucleus—the new macronucleus, to influence epigenetic DNA elimination in conjugating cells (Figure 4, right; Mochizuki & Gorovsky, 2005). Many important questions remain: (1) How are dsRNAs are imported into nuclei? (2) Do the same dsRNAs that participate in chromatin alteration also participate in posttranscriptional events? (3) How do cells distinguish between dsRNAs that function in the cytoplasm and those that function in the nucleus?

## SUMMARY AND PERSPECTIVES

The subcellular dynamics of small RNAs encoded by the nucleus are unexpectedly complex. Some small RNAs that function solely in the nucleus, such as snRNAs, are processed in the cytoplasm before they return to the nucleus (Figure 1). It is not clear whether

this seemingly complicated life style for snRNAs is conserved among all eukaryotes, but some recent tantalizing evidence for budding yeast suggest that it may be. It is also not clear whether other types of small RNAs that function in the nucleus sample the cytoplasm during their biogenesis, but new studies in budding yeast for telomerase RNA (Figure 2), and in vertebrate cells for snoRNAs, suggest that these RNAs may also sample the cytoplasm. In budding yeast tRNAs undergo very intricate cellular dynamics (Figure 3). Pre-tRNAs exit the nucleus and complete their processing in the cytoplasm. Some or all of the processed cytoplasmic tRNAs move retrograde back to the nucleus and likely return once again to the cytoplasm via a re-export process. Whether the retrograde process is regulated is unknown; it is also unknown whether it is conserved among other eukaryotes. Small dsRNAs may also have complex nucleus/cytoplasmic dynamics. There is one documented case for members of this group (scnRNAs) traveling from one nucleus to the cytoplasm and then to a second and third nucleus (Figure 4); it is unknown whether other dsRNAs sample the cytoplasm before functioning in the nucleus. Future experiments must address the details of the mechanism(s) for each category of traveling RNA, determine whether the travels are conserved among eukaryotes, and elucidate the functional significance that such seemingly complicated cell biology serves.

## ACKNOWLEDGMENTS

I am grateful for scientific conversations with M. Gorovsky, E. Phizicky, P. Silicano, and M. Terns. I thank members of my laboratory, R. Hurto, A. Murthi, H. Shaheen, D. Stanford, K. Stauffer, and M. Whitney for scientific feedback and comments on the manuscript. I also thank E. Phizicky and J. E. Hopper for comments on the manuscript. The work in my laboratory was supported by a grant (GM27930) from the NIH.

## REFERENCES

Aebi, M., Clark, M.W., Vijayraghavan, U., and Abelson, J. 1990. A yeast mutant, *PRP20*, altered in mRNA metabolism and maintenance of the nuclear structure, is defective in a gene homologous to the human gene *RCC1* which is involved in the control of chromosome condensation. *Mol Gen Genet* 224:72–80.

Arts, G.J., Fornerod, M., and Mattaj, I.W. 1998a. Identification of a nuclear export receptor for tRNA. *Curr Biol* 8:305–314.

Arts, G.J., Kuersten, S., Romby, P., Ehresmann, B., and Mattaj, I.W. 1998b. The role of exportin-t in selective nuclear export of mature tRNAs. *EMBO J* 17:7430–7441.

Azad, A.K., Stanford, D.R., Sarkar, S., and Hopper, A.K. 2001. Role of nuclear pools of aminoacyl-tRNA synthetases in tRNA nuclear export. *Mol Biol Cell* 12:1381–1392.

Bagga, S., Bracht, J., Massirer, K., Holtz, J., Eachus, R., and Pasquinelli, A.E. 2005. Regulation by let-7 and lin-4 miRNAs results in target mRNA degradation. *Cell* 122:553–568.

Bartel, D.P. 2004. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116:281–297.

Baserga, S.J., Gilmore-Hebert, M., and Yang, X.W. 1992. Distinct molecular signals for nuclear import of the nucleolar snRNA, U3. *Genes Dev* 6:1120–1130.

Bernstein, E. and Allis, C.D. 2005. RNA meets chromatin. *Genes Dev* 19:1635–1655.

Bertrand, E., Houser-Scott, F., Kendall, A., Singer, R.H., and Engelke, D.R. 1998. Nucleolar localization of early tRNA processing. *Genes Dev* 12:2463–2468.

Bohnsack, M.T., Czaplinski, K., and Görlich, D. 2004. Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs. *RNA* 10:185–191.

Bohnsack, M.T., Regener, K., Schwappach, B., Saffrich, R., Paraskeva, E., Hartmann, E., and Görlich, D. 2002. Exp5 exports eEF1A via tRNA from nuclei and synergizes with other transport pathways to confine translation to the cytoplasm. *EMBO J* 21:6205–6215.

Boulon, S., Verheggen, C., Jady, B.E., Girard, C., Pescia, C., Paul, C., Ospina, J.K., Kiss, T., Matera, A.G., Bordonne, R., and Bertrand, E. 2004. PHAX and CRM1 are required sequentially to transport U3 snoRNA to nucleoli. *Mol Cell* 16:777–787.

Calado, A., Treichel, N., Muller, E.C., Otto, A., and Kutay, U. 2002. Exportin-5-mediated nuclear export of eukaryotic elongation factor 1A and tRNA. *EMBO J* 21:6216–6224.

Cech, T.R. 2004. Beginning to understand the end of the chromosome. *Cell* 116:273–279.

Cheng, Y., Dahlberg, J.E., and Lund, E. 1995. Diverse effects of the guanine nucleotide exchange factor RCC1 on RNA transport. *Science* 267:1807–1810.

Collins, K. 1999. Ciliate telomerase biochemistry. *Annu Rev Biochem* 68:187–218.

Cronshaw, J.M., Krutchinsky, A.N., Zhang, W., Chait, B.T., and Matunis, M.J. 2002. Proteomic analysis of the mammalian nuclear pore complex. *J Cell Biol* 158:915–927.

Dahlberg, J.E., Lund, E., and Goodwin, E.B. 2003. Nuclear translation: what is the evidence? *RNA* 9:1–8.

De Robertis, E.M., Black, P., and Nishikura, K. 1981. Intranuclear location of the tRNA splicing enzymes. *Cell* 23:89–93.

Dingermann, T., Werner, H., Schutz, A., Zundorf, I., Nerke, K., Knecht, D., and Marschalek, R. 1992. Establishment of a system for conditional gene expression using an inducible tRNA suppressor gene. *Mol Cell Biol* 12:4038–4045.

Dutcher, S.K. 1981. Internuclear transfer of genetic information in *kar1-1/KAR1* heterokaryons in *Saccharomyces cerevisiae*. *Mol Cell Biol* 1:245–253.

Feng, W., and Hopper, A.K. 2002. A Los1p-independent pathway for nuclear export of intronless tRNAs in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci USA* 99:5412–5417.

Ferrezzuelo, F., Steiner, B., Aldea, M., and Fitcher, B. 2002. Biogenesis of yeast telomerase depends on the importin Mtr10. *Mol Cell Biol* 22:6046–6055.

Fischer, U. and Lührmann, R. 1990. An essential signaling role for the m<sup>3</sup>G cap in the transport of U1 snRNP to the nucleus. *Science* 249:786–790.

Flach, J., Bossie, M., Vogel, J., Corbett, A., Jinks, T., Willins, D.A., and Silver, P.A. 1994. A yeast RNA-binding protein shuttles between the nucleus and the cytoplasm. *Mol Cell Biol* 14:8399–8407.

Görlich, D. and Kutay, U. 1999. Transport between the cell nucleus and the cytoplasm. *Annu Rev Cell Dev Biol* 15:607–660.

Grosshans, H., Hurt, E., and Simos, G. 2000. An aminoacylation-dependent nuclear tRNA export pathway in yeast. *Genes Dev* 14:830–840.

- Gu, W., Jackman, J.E., Lohan, A.J., Gray, M.W., and Phizicky, E.M. 2003. tRNA<sup>His</sup> maturation: An essential yeast protein catalyzes addition of a guanine nucleotide to the 5' end of tRNA<sup>His</sup>. *Genes Dev* 17:2889–2901.
- Gwizdek, C., Ossareh-Nazari, B., Brownawell, A.M., Evers, S., Macara, I.G., and Dargemont, C. 2004. Minihelix-containing RNAs mediate exportin-5-dependent nuclear export of the double-stranded RNA-binding protein ILF3. *J Biol Chem* 279:884–891.
- Haeusler, R.A. and Engelke, D.R. 2004. Genome organization in three dimensions: thinking outside the line. *Cell Cycle* 3:273–275.
- Hamm, J. and Mattaj, I.W. 1990. Monomethylated cap structures facilitate RNA export from the nucleus. *Cell* 63:109–118.
- Hausmann, S. and Shuman, S. 2005. Specificity and mechanism of RNA cap guanine-N<sup>2</sup> methyltransferase (Tgs1). *J Biol Chem* 280:4021–4024.
- Hellmuth, K., Lau, D.M., Bischoff, F.R., Kunzler, M., Hurt, E., and Simos, G. 1998. Yeast Los1p has properties of an exportin-like nucleocytoplasmic transport factor for tRNA. *Mol Cell Biol* 18:6374–6386.
- Hiley, S.L., Babak, T., and Hughes, T.R. 2005. Global analysis of yeast RNA processing identifies new targets of RNase III and uncovers a link between tRNA 5' end processing and tRNA splicing. *Nucleic Acids Res* 33:3048–3056.
- Hopper, A.K., Banks, F., and Evangelides, V. 1978. A yeast mutant which accumulates precursor tRNAs. *Cell* 14:211–219.
- Hopper, A.K. and Phizicky, E.M. 2003. tRNA transfers to the limelight. *Genes Dev* 17:162–180.
- Hopper, A.K., Schultz, L.D., and Shapiro, R.A. 1980. Processing of intervening sequences: a new yeast mutant which fails to excise intervening sequences from precursor tRNAs. *Cell* 19:741–751.
- Hopper, A.K., Traglia, H.M., and Dunst, R.W. 1990. The yeast *RNA1* gene product necessary for RNA processing is located in the cytosol and apparently excluded from the nucleus. *J Cell Biol* 111:309–321.
- Huber, J., Cronshagen, U., Kadokura, M., Marshallsay, C., Wada, T., Sekine, M., and Lührmann, R. 1998. Snurportin1, an m<sup>3</sup>G-cap-specific nuclear import receptor with a novel domain structure. *EMBO J* 17:4114–4126.
- Huh, W.K., Falvo, J.V., Gerke, L.C., Carroll, A.S., Howson, R.W., Weissman, J.S., and O'Shea, E.K. 2003. Global analysis of protein localization in budding yeast. *Nature* 425:686–691.
- Hunter, C.A., Aukerman, M.J., Sun, H., Fokina, M., and Poethig, R.S. 2003. PAUSED encodes the Arabidopsis exportin-t ortholog. *Plant Physiol* 132:2135–2143.
- Hurt, D.J., Wang, S.S., Lin, Y.H., and Hopper, A.K. 1987. Cloning and characterization of *LOS1*, a *Saccharomyces cerevisiae* gene that affects tRNA splicing. *Mol Cell Biol* 7:1208–1216.
- Iborra, F.J., Jackson, D.A., and Cook, P.R. 2001. Coupled transcription and translation within nuclei of mammalian cells. *Science* 293:1139–1142.
- Izaurralde, E., Kutay, U., von Kobbe, C., Mattaj, I.W., and Görlich, D. 1997. The asymmetric distribution of the constituents of the Ran system is essential for transport into and out of the nucleus. *EMBO J* 16:6535–6547.
- Izaurralde, E., Lewis, J., Gamberi, C., Jarmolowski, A., McGuigan, C., and Mattaj, I.W. 1995. A cap-binding protein complex mediating U snRNA export. *Nature* 376:709–712.
- Kadaba, S., Krueger, A., Trice, T., Krecic, A.M., Hinnebusch, A.G., and Anderson, J. 2004. Nuclear surveillance and degradation of hypomethylated initiator tRNA<sup>Met</sup> in *S. cerevisiae*. *Genes Dev* 18:1227–1240.
- Kadowaki, T., Chen, S., Hitomi, M., Jacobs, E., Kumagai, C., Liang, S., Schreiber, R., Singleton, D., Wisniewska, J., and Tartakoff, A.M. 1994. Isolation and characterization of *Saccharomyces cerevisiae* mRNA transport-defective (mtr) mutants. *J Cell Biol* 126:649–659.
- Kalab, P., Weis, K., and Heald, R. 2002. Visualization of a Ran-GTP gradient in interphase and mitotic *Xenopus* egg extracts. *Science* 295:2452–2456.
- Kato, H., Goto, D.B., Martienssen, R.A., Urano, T., Furukawa, K., and Murakami, Y. 2005. RNA polymerase II is required for RNAi-dependent heterochromatin assembly. *Science* 309:467–469.
- Kiss, T. 2002. Small nucleolar RNAs: an abundant group of noncoding RNAs with diverse cellular functions. *Cell* 109:145–148.
- Kiss, T. 2004. Biogenesis of small nuclear RNPs. *J Cell Sci* 117:5949–5951.
- Kuersten, S., Arts, G.J., Walther, T.C., Englmeier, L., and Mattaj, I.W. 2002. Steady-state nuclear localization of exportin-t involves RanGTP binding and two distinct nuclear pore complex interaction domains. *Mol Cell Biol* 22:5708–5720.
- Kutay, U., Lipowsky, G., Izaurralde, E., Bischoff, F.R., Schwarzmaier, P., Hartmann, E., and Görlich, D. 1998. Identification of a tRNA-specific nuclear export receptor. *Mol Cell* 1:359–369.
- Lamond, A.I. and Spector, D.L. 2003. Nuclear speckles: a model for nuclear organelles. *Nat Rev Mol Cell Biol* 4:605–612.
- Leary, D.J., Terns, M.P., and Huang, S. 2004. Components of U3 snoRNA-containing complexes shuttle between nuclei and the cytoplasm and differentially localize in nucleoli: implications for assembly and function. *Mol Biol Cell* 15:281–293.
- Lee, Y., Ahn, C., Han, J., Choi, H., Kim, J., Yim, J., Lee, J., Provost, P., Radmark, O., Kim, S., and Kim, V.N. 2003. The nuclear RNase III Drosha initiates microRNA processing. *Nature* 425:415–419.
- Lendvai, T.S., Morris, D.K., Sah, J., Balasubramanian, B., and Lundblad, V. 1996. Senescence mutants of *Saccharomyces cerevisiae* with a defect in telomere replication identify three additional EST genes. *Genetics* 144:1399–1412.
- Li, J. and Chen, X. 2003. PAUSED, a putative exportin-t, acts pleiotropically in *Arabidopsis* development but is dispensable for viability. *Plant Physiol* 132:1913–1924.
- Lingner, J., Hughes, T.R., Shevchenko, A., Mann, M., Lundblad, V., and Cech, T.R. 1997. Reverse transcriptase motifs in the catalytic subunit of telomerase. *Science* 276:561–567.
- Lipowsky, G., Bischoff, F.R., Izaurralde, E., Kutay, U., Schafer, S., Gross, H.J., Beier, H., and Görlich, D. 1999. Coordination of tRNA nuclear export with processing of tRNA. *RNA* 5:539–549.
- Liu, J., Valencia-Sanchez, M.A., Hannon, G.J., and Parker, R. 2005. MicroRNA-dependent localization of targeted mRNAs to mammalian P-bodies. *Nat Cell Biol* 7:719–723.
- Lukowiak, A.A., Narayanan, A., Li, Z.H., Terns, R.M., and Terns, M.P. 2001. The snoRNA domain of vertebrate telomerase RNA functions to localize the RNA within the nucleus. *RNA* 7:1833–1844.
- Lund, E. and Dahlberg, J.E. 1998. Proofreading and aminoacylation of tRNAs before export from the nucleus. *Science* 282:2082–2085.
- Lund, E., Guttinger, S., Calado, A., Dahlberg, J.E., and Kutay, U. 2004. Nuclear export of microRNA precursors. *Science* 303:95–98.
- Martienssen, R.A., Zaratiegui, M., and Goto, D.B. 2005. RNA interference and heterochromatin in the fission yeast *Schizosaccharomyces pombe*. *Trends Genet* 21:450–456.
- Maser, R.S. and DePinho, R.A. 2002. Keeping telomerase in its place. *Nat Med* 8:934–936.
- Matera, A.G. 1999. Nuclear bodies: multifaceted subdomains of the interchromatin space. *Trends Cell Biol* 9:302–309.
- Matzke, M.A. and Birchler, J.A. 2005. RNAi-mediated pathways in the nucleus. *Nat Rev Genet* 6:24–35.
- Michaud, N. and Goldfarb, D. 1992. Microinjected U snRNAs are imported to oocyte nuclei via the nuclear pore complex by three distinguishable targeting pathways. *J Cell Biol* 116:851–861.
- Mitchell, J.R., Cheng, J., and Collins, K. 1999. A box H/ACA small nucleolar RNA-like domain at the human telomerase RNA 3' end. *Mol Cell Biol* 19:567–576.
- Mochizuki, K. and Gorovsky, M.A. 2005. A Dicer-like protein in *Tetrahymena* has distinct functions in genome rearrangement, chromosome segregation, and meiotic prophase. *Genes Dev* 19:77–89.
- Mouaikel, J., Verheggen, C., Bertrand, E., Tazi, J., and Bordonne, R. 2002. Hypermethylation of the cap structure of both yeast snRNAs and snoRNAs requires a conserved methyltransferase that is localized to the nucleolus. *Mol Cell* 9:891–901.

- Murchison, E.P. and Hannon, G.J. 2004. miRNAs on the move: miRNA biogenesis and the RNAi machinery. *Curr Opin Cell Biol* 16:223–229.
- Murphy, M.W., Olson, B.L., and Siliciano, P.G. 2004. The yeast splicing factor Prp40p contains functional leucine-rich nuclear export signals that are essential for splicing. *Genetics* 166:53–65.
- Narayanan, A., Eifert, J., Marfatia, K.A., Macara, I.G., Corbett, A.H., Terns, R.M., and Terns, M.P. 2003. Nuclear RanGTP is not required for targeting small nucleolar RNAs to the nucleolus. *J Cell Sci* 116:177–186.
- Narayanan, A., Lukowiak, A., Jady, B.E., Dragon, F., Kiss, T., Terns, R.M., and Terns, M.P. 1999. Nucleolar localization signals of box H/ACA small nucleolar RNAs. *EMBO J* 18:5120–5130.
- Narayanan, U., Achsel, T., Lührmann, R., and Matera, A.G. 2004. Coupled in vitro import of U snRNPs and SMN, the spinal muscular atrophy protein. *Mol Cell* 16:223–234.
- Nathanson, L. and Deutscher, M.P. 2000. Active aminoacyl-tRNA synthetases are present in nuclei as a high molecular weight multienzyme complex. *J Biol Chem* 275:31559–31562.
- Noma, K., Sugiyama, T., Cam, H., Verdel, A., Zofall, M., Jia, S., Moazed, D., and Grewal, S.I. 2004. RITS acts in cis to promote RNA interference-mediated transcriptional and post-transcriptional silencing. *Nat Genet* 36:1174–1180.
- O'Connor, J.P. and Peebles, C.L. 1991. In vivo pre-tRNA processing in *Saccharomyces cerevisiae*. *Mol Cell Biol* 11:425–439.
- Ohno, M., Segref, A., Bachi, A., Wilm, M., and Mattaj, I.W. 2000. PHAX, a mediator of U snRNA nuclear export whose activity is regulated by phosphorylation. *Cell* 101:187–198.
- Ohtsubo, M., Yoshida, T., Seino, H., Nishitani, H., Clark, K.L., Sprague, G.F., Jr., Frasch, M., and Nishimoto, T. 1991. Mutation of the hamster cell cycle gene RCC1 is complemented by the homologous genes of *Drosophila* and *S. cerevisiae*. *EMBO J* 10:1265–1273.
- Olson, B.L. and Siliciano, P.G. 2003. A diverse set of nuclear RNAs transfer between nuclei of yeast heterokaryons. *Yeast* 20:893–903.
- Park, M.Y., Wu, G., Gonzalez-Sulser, A., Vaucheret, H., and Poethig, R.S. 2005. Nuclear processing and export of microRNAs in *Arabidopsis*. *Proc Natl Acad Sci USA* 102:3691–3696.
- Pasquinelli, A.E., Hunter, S., and Bracht, J. 2005. MicroRNAs: a developing story. *Curr Opin Genet Dev* 15:200–205.
- Paushkin, S.V., Patel, M., Furia, B.S., Peltz, S.W., and Trotta, C.R. 2004. Identification of a human endonuclease complex reveals a link between tRNA splicing and pre-mRNA 3' end formation. *Cell* 117:311–321.
- Peculis, B.A. 2001. snoRNA nuclear import and potential for cotranscriptional function in pre-rRNA processing. *RNA* 7:207–219.
- Pillai, R.S., Bhattacharyya, S.N., Artus, C.G., Zoller, T., Cougot, N., Basyuk, E., Bertrand, E., and Filipowicz, W. 2005. Inhibition of translational initiation by let-7 MicroRNA in Human Cells. *Science* 309:1573–1576.
- Popenko, V.I., Ivanova, J.L., Cherny, N.E., Filonenko, V.V., Beresten, S.F., Wolfson, A.D., and Kisselev, L.L. 1994. Compartmentalization of certain components of the protein synthesis apparatus in mammalian cells. *Eur J Cell Biol* 65:60–69.
- Rodriguez, M.S., Dargemont, C., and Stutz, F. 2004. Nuclear export of RNA. *Biol Cell* 96:639–655.
- Rose, A.M., Belford, H.G., Shen, W.C., Greer, C.L., Hopper, A.K., and Martin, N.C. 1995. Location of N<sup>2</sup>, N<sup>2</sup>-dimethylguanosine-specific tRNA methyltransferase. *Biochimie* 77:45–53.
- Rose, A.M., Joyce, P.B., Hopper, A.K., and Martin, N.C. 1992. Separate information required for nuclear and subnuclear localization: additional complexity in localizing an enzyme shared by mitochondria and nuclei. *Mol Cell Biol* 12:5652–5658.
- Rout, M.P., Aitchison, J.D., Suprapto, A., Hjertaas, K., Zhao, Y., and Chait, B.T. 2000. The yeast nuclear pore complex: composition, architecture, and transport mechanism. *J Cell Biol* 148:635–651.
- Rozenski, J., Crain, P.F., and McCloskey, J.A. 1999. The RNA Modification Database: 1999 update. *Nucleic Acids Res* 27:196–197.
- Sarkar, S., Azad, A.K., and Hopper, A.K. 1999. Nuclear tRNA aminoacylation and its role in nuclear export of endogenous tRNAs in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci USA* 96:14366–14371.
- Sarkar, S. and Hopper, A.K. 1998. tRNA nuclear export in *Saccharomyces cerevisiae*: in situ hybridization analysis. *Mol Biol Cell* 9:3041–3055.
- Schramke, V., Sheedy, D.M., Denli, A.M., Bonila, C., Ekwall, K., Hannon, G.J., and Allshire, R.C. 2005. RNA-interference-directed chromatin modification coupled to RNA polymerase II transcription. *Nature* 435:1275–1279.
- Sen, G.L. and Blau, H.M. 2005. Argonaute 2/RISC resides in sites of mammalian mRNA decay known as cytoplasmic bodies. *Nat Cell Biol* 7:633–636.
- Senger, B., Simos, G., Bischoff, F.R., Podtelejnikov, A., Mann, M., and Hurt, E. 1998. Mtr10p functions as a nuclear import receptor for the mRNA-binding protein Npl3p. *EMBO J* 17:2196–2207.
- Seto, A.G., Zaug, A.J., Sobel, S.G., Wolin, S.L., and Cech, T.R. 1999. *Saccharomyces cerevisiae* telomerase is an Sm small nuclear ribonucleoprotein particle. *Nature* 401:177–180.
- Shaheen, H.H. and Hopper, A.K. 2005. Retrograde movement of tRNAs from the cytoplasm to the nucleus in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci USA* 102:12290–12295.
- Sharma, K., Fabre, E., Tekotte, H., Hurt, E.C., and Tollervey, D. 1996. Yeast nucleoporin mutants are defective in pre-tRNA splicing. *Mol Cell Biol* 16:294–301.
- Shimba, S., Buckley, B., Reddy, R., Kiss, T., and Filipowicz, W. 1992. Cap structure of U3 small nucleolar RNA in animal and plant cells is different. gamma-Monomethyl phosphate cap structure in plant RNA. *J Biol Chem* 267:13772–13777.
- Shulga, N., Mosammaparast, N., Wozniak, R., and Goldfarb, D.S. 2000. Yeast nucleoporins involved in passive nuclear envelope permeability. *J Cell Biol* 149:1027–1038.
- Sigurdson, D.C., Gaarder, M.E., and Livingston, D.M. 1981. Characterization of the transmission during cytoductant formation of the 2 micrometers DNA plasmid from *Saccharomyces*. *Mol Gen Genet* 183:59–65.
- Singh, R. and Reddy, R. 1989. Gamma-monomethyl phosphate: a cap structure in spliceosomal U6 small nuclear RNA. *Proc Natl Acad Sci USA* 86:8280–8283.
- Sontheimer, E.J. 2005. Assembly and function of RNA silencing complexes. *Nat Rev Mol Cell Biol* 6:127–138.
- Speckmann, W., Narayanan, A., Terns, R., and Terns, M.P. 1999. Nuclear retention elements of U3 small nucleolar RNA. *Mol Cell Biol* 19:8412–8421.
- Steiner-Mosonyi, M. and Mangroo, D. 2004. The nuclear tRNA aminoacylation-dependent pathway may be the principal route used to export tRNA from the nucleus in *Saccharomyces cerevisiae*. *Biochem J* 378:809–816.
- Takano, A., Endo, T. and Yoshihisa, T. 2005. tRNA actively shuttles between the nucleus and cytosol in yeast. *Science* 309:140–142.
- Tang, G. 2005. siRNA and miRNA: an insight into RISCs. *Trends Biochem Sci* 30:106–114.
- Tarn, W.Y. and Steitz, J.A. 1997. Pre-mRNA splicing: the discovery of a new spliceosome doubles the challenge. *Trends Biochem Sci* 22:132–137.
- Teixeira, D., Sheth, U., Valencia-Sanchez, M.A., Brengues, M., and Parker, R. 2005. Processing bodies require RNA for assembly and contain nontranslating mRNAs. *RNA* 11:371–382.
- Teixeira, M.T., Forstemann, K., Gasser, S.M., and Lingner, J. 2002. Intracellular trafficking of yeast telomerase components. *EMBO Rep* 3:652–659.
- Terns, M.P. and Dahlberg, J.E. 1994. Retention and 5' cap trimethylation of U3 snRNA in the nucleus. *Science* 264:959–961.
- Terns, M.P., Dahlberg, J.E., and Lund, E. 1993. Multiple cis-acting signals for export of pre-U1 snRNA from the nucleus. *Genes Dev* 7:1898–1908.
- Terns, M.P., Grimm, C., Lund, E., and Dahlberg, J.E. 1995. A common maturation pathway for small nucleolar RNAs. *EMBO J* 14:4860–4871.

- Terns, M.P. and Terns, R.M. 2001. Macromolecular complexes:SMN—the master assembler. *Curr Biol* 11:R862–864.
- Thompson, M., Haeusler, R.A., Good, P.D., and Engelke, D.R. 2003. Nucleolar clustering of dispersed tRNA genes. *Science* 302:1399–1401.
- Tolerico, L.H., Benko, A.L., Aris, J.P., Stanford, D.R., Martin, N.C., and Hopper, A.K. 1999. *Saccharomyces cerevisiae* Mod5p-II contains sequences antagonistic for nuclear and cytosolic locations. *Genetics* 151:57–75.
- Tomari, Y. and Zamore, P.D. 2005. Perspective: machines for RNAi. *Genes Dev* 19:517–529.
- Tycowski, K.T., Shu, M.D., and Steitz, J.A. 1996. A mammalian gene with introns instead of exons generating stable RNA products. *Nature* 379:464–466.
- van Hoof, A., Lennertz, P., and Parker, R. 2000. Yeast exosome mutants accumulate 3'-extended polyadenylated forms of U4 small nuclear RNA and small nucleolar RNAs. *Mol Cell Biol* 20:441–452.
- Vankan, P., McGuigan, C., and Mattaj, I.W. 1990. Domains of U4 and U6 snRNAs required for snRNP assembly and splicing complementation in *Xenopus* oocytes. *EMBO J* 9:3397–3404.
- Watkins, N.J., Lemm, I., Ingelfinger, D., Schneider, C., Hossbach, M., Urlaub, H., and Luhrmann, R. 2004. Assembly and maturation of the U3 snoRNP in the nucleoplasm in a large dynamic multiprotein complex. *Mol Cell* 16:789–798.
- Wu, P., Brockenbrough, J.S., Metcalfe, A.C., Chen, S., and Aris, J.P. 1998. Nop5p is a small nucleolar ribonucleoprotein component required for pre-18 S rRNA processing in yeast. *J Biol Chem* 273:16453–16463.
- Xiao, S., Scott, F., Fierke, C.A., and Engelke, D.R. 2002. Eukaryotic ribonuclease P: a plurality of ribonucleoprotein enzymes. *Annu Rev Biochem* 71:165–189.
- Yi, R., Qin, Y., Macara, I.G., and Cullen, B.R. 2003. Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes Dev* 17:3011–3016.
- Yong, J., Wan, L., and Dreyfuss, G. 2004. Why do cells need an assembly machine for RNA-protein complexes? *Trends Cell Biol* 14:226–232.
- Yoshida, K. and Blobel, G. 2001. The karyopherin Kap142p/Msn5p mediates nuclear import and nuclear export of different cargo proteins. *J Cell Biol* 152:729–740.
- Yoshihisa, T., Yunoki-Esaki, K., Ohshima, C., Tanaka, N., and Endo, T. 2003. Possibility of cytoplasmic pre-tRNA splicing: the yeast tRNA splicing endonuclease mainly localizes on the mitochondria. *Mol Biol Cell* 14:3266–3279.
- Zeng, Y. and Cullen, B.R. 2005. Efficient processing of primary microRNA hairpins by Drosha requires flanking nonstructured RNA sequences. *J Biol Chem* 280:27595–27603.

Editor: Marvin Wickens